arXiv:1609.01695v1 [math.SP] 6 Sep 2016

A trace formula for the index of B-Fredholm operators

M. Berkani

Abstract

In this paper we define B-Fredholm elements in a Banach algebra A modulo an ideal J of A. When a trace function is given on the ideal J, it generate an index for B-Fredholm elements. In the case of a B-Fredholm operator T acting on a Banach space, we prove that its usual index $ind(T)$ is equal to the trace of the commutator $[T, T_0]$, where T_0 is a Drazin inverse of T modulo the ideal of finite rank operators, extending a Fedosov's trace formula for Fredholm operators [8]. In the case of a semi-simple Banach algebra, we prove a punctured neighborhood theorem for the index.

1 Introduction

Let X be a Banach space and let $L(X)$ be the Banach algebra of bounded linear operators acting on X. In [3], we have introduced the class of linear bounded B-Fredholm operators. If $F_0(X)$ is the ideal of finite rank operators in $L(X)$ and $\pi : L(X) \longrightarrow A$ is the canonical projection, from $L(X)$ onto the quotient algebra $A = L(X)/F_0(X)$, it is well known by the Atkinson's theorem [2, Theorem 0.2.2, p.4], that $T \in L(X)$ is a Fredholm operator if and only if its projection $\pi(T)$ in the algebra A is invertible. Similarly, in the following result, we have established an Atkinson-type theorem for B-Fredholm operators.

THEOREM 1.1. *[6, Theorem 3.4]: Let $T \in L(X)$. Then T is a B-Fredholm operator if and only if $\pi(T)$ is Drazin invertible in the algebra $L(X)/F_0(X)$.*

2010 Mathematics Subject Classification: Primary 47A53, 46H05

Key words and phrases: B-Fredholm, Banach algebra, index,trace

Motivated by this result, we define in this paper B-Fredholm elements in a semi-prime unital Banach algebra A modulo an ideal J of A. Recall that a Banach algebra A is called semi-prime if for $u \in A, uxu = 0$, for all $x \in A$ implies that $u = 0$.
An element $a \neq 0$ in a semiprime Banach algebra A is called of rank one if there exists a linear functional f_a on A such that $axa = f_a(x)a$ for all $x \in A$.

Definition 1.2. Let A be unital semi-prime Banach algebra and let J be an ideal of A, and $\pi : A \to A/J$ the canonical projection. An element $a \in A$ is called a B-Fredholm element of A modulo the ideal J if its image $\pi(a)$ is Drazin invertible in the quotient algebra A/J.

In a recent work [9], the authors, gave in [9, Definiton 2.3] a definition of B-Fredholm elements in Banach algebras. However, their definition does not englobe the class of B-Fredholm operators, since the algebra $L(X)/F_0(X)$ is not a Banach algebra. That's why in our definition, we consider general algebras, not necessarily being Banach algebras, so it includes also the case of the algebra $L(X)/F_0(X)$. While an extensive study of B-Fredholm elements in a Banach algebras modulo an ideal J is being done in [7], we focus our attention here on the properties of the index of such elements.
Recently in [10], the authors studied Fredholm elements in a semi-prime Banach algebra modulo an ideal. When a trace function is given on the ideal considered, they define also the index of a Fredholm element using a trace formula [10, Definition 3.3]. This definition coincides with the usual definition of the index for a Fredholm operator acting on a Banach space.
In the second section of this paper, following the same approach as in [10], when a trace function is defined on the ideal J considered, we will define an index for B-Fredholm elements of the Banach algebra A modulo the ideal J. Then in the case of B-Fredholm operator T acting on a Banach space X, we prove our main result announced in the abstract, that is the usual index $ind(T)$ is equal to $\tau([T, T_0])$ where T_0 is a Drazin inverse of T modulo the ideal of finite rank operators.
In the third section, we give a punctured neighborhood theorem for the index B-Fredholm elements in a semi-simple Banach algebra. We will also establish a logarithmic rule of the index, for two commuting prime B-Fredholm elements.

2 Trace and Index

In this section, A will be a semi-prime, complex and unital Banach algebra. We consider here B-Fredholm elements in A modulo an ideal J, on which a trace

function is given. Then we extend the definition of the index [10, Definition 3.3] to the class of B-Fredholm elements of A modulo J. We will show also that the new definition of the index coincides with the usual one in the case of B-Fredholm operator acting on a Banach space given in [3, Definition 2.3].

DEFINITION 2.1. [10, 2.1., p.283] Let J be an ideal in a Banach algebra A. A function $\tau : J \to \mathbb{C}$, is called a trace on J if :

1)$\tau(p) = 1$ if $p \in J$ is an idempotent, that is $p^2 = p$, and p of rank one,
2) $\tau(a + b) = \tau(a) + \tau(b)$, for all $a, b \in J$,
3) $\tau(\alpha a) = \alpha\tau(a)$, forall $\alpha \in \mathbb{C}$ and $a \in J$,
4) $\tau(ab) = \tau(ba)$, for all $a \in J$ and $b \in A$.

DEFINITION 2.2. Let τ be a trace on an ideal J of a Banach algebra A. The index of a B-Fredholm element $a \in A$ is defined by:

$$\mathbf{i}(a) = \tau(aa_0 - a_0a) = \tau([a, a_0]),$$

where a_0 is a Drazin inverse of a modulo the ideal I.

THEOREM 2.3. *The index of a B-Fredholm element $a \in A$ is well defined and is independant of the Drazin inverse a_0 of a modulo the ideal J.*

Proof. Let $a \in A$ be a B-Fredholm element in A modulo J. Then $\pi(a)$ is Drazin invertible in A/J. If $\pi(a_0)$ is the Drazin inverse of $\pi(a)$, then $\pi(a)\pi(a_0) = \pi(a_0)\pi(a)$. Hence $aa_0 - a_0a \in J$. If $\pi(a_0) = \pi(a_0')$, then $a_0' - a_0 \in J$ and $aa_0' - a_0'a - (aa_0 - a_0a) = a(a_0' - a_0) - (a_0' - a_0)a$. Since $(a_0' - a_0) \in J$, using the property 4) of the trace, we obtain $\tau(a(a_0' - a_0)) = \tau((a_0' - a_0)a)$. So the index is independent of the choice of the representative a_0.

REMARK 2.4. Clearly the definition of the index of B-Fredholm elements, extends [10, Definition 3.3], given for Fredholm elements in Banach algebras, because the inverse modulo J of a Fredholm element $a \in A$ is also its Drazin inverse modulo J.

Now we prove in the next theorem, that the index defined here for B-Fredholm elements in Banach algebras is equal to the usual index of B-Fredholm operators acting on a Banach space. Here the ideal I is the ideal of finite rank operators acting on X, and the trace τ is the trace that is defined for finite rank operators as in [10, Example 2.1]. If $T \in I$ and $\{x_1, ..., x_n\}$ is a basis of its image $R(T)$, then $T = \Sigma_{i=1}^n x_i' \bigotimes x_i$, where $x_1', ..., x_n'$ are continuous linear functionals on X such that $T(x) = \Sigma_{i=1}^n x_i'(x)x_i$, for all $x \in X$. The trace of T is then defined by:

$$\tau(T) = \Sigma_{i=1}^n x_i'(x_i).$$

THEOREM 2.5. *Let X be a Banach space and let $T \in B(X)$ be a B-Fredholm operator. Then its usual index $ind(T)$ is equal to the trace $\tau([T,S])$ where S is a Drazin inverse of T modulo the ideal of finite rank operators.*

Proof. Let T be a B-Fredholm operator, then from [3, Theorem 2.7] there exist two closed subspaces M and N of X such that $X = M \oplus N$ and:
i) $T(N) \subset N$ and $T_{|N}$ is a nilpotent operator,
ii) $T(M) \subset M$ and $T_{|M}$ is a Fredholm operator.
Let $p \in \mathbb{N}$, such that $(T_{|N})^p = 0$. Then $T^p = (T_{|M})^p \oplus 0$. From [3, Theorem 2.7], T^p is a B-Fredholm operator and from [3, Proposition 2.1] $ind(T^p) = p.ind(T) = p\, ind(T_{|M})$. Let T_0 be an inverse of $T_{|M}$ modulo the finite rank operators on M and $S = T_0 \oplus 0$. Then $S = T_0 \oplus 0$ is a Drazin inverse of T and $S^p = (T_0)^p \oplus 0$ is a Drazin inverse of T^p modulo the finite rank operators on X. Moreover $T^pS^p - S^pT^p = [(T_{|M})^pT_0^p - T_0^p(T_{|M})^p] \oplus 0$.
We observe that $(T_{|M})^pT_0^p - T_0^p(T_{|M})^p$ is of finite rank. So $T^pS^p - S^pT^p$ is also of finite rank. Since T_0^p is an inverse of the Fredholm operator $(T_{|M})^p$ modulo the finite rank on $T_{|M}$, then from [10, Example 3.2],we have $\tau(T^pS^p - S^pT^p) = \tau((T_{|M})^pT_0^p - T_0^p(T_{|M})^p) = ind((T_{|M})^p) = p\, ind(T_{|M})) = p\ ind(T)$. Here τ, is the function trace defined on finite rank operators acting on a Banach space as in [10, Examlpe 2.1].
On another side, by [10, Proposition 3.5] we have $\tau((T_{|M})^pT_0^p - T_0^p(T_{|M})^p) = \mathbf{i}((T_{|M})^p) = p\, \mathbf{i}(T_{|M})$ and so $ind(T) = \mathbf{i}(T_{|M})$. Since, by definition of the index $\mathbf{i}$, we have $\mathbf{i}(T) = \tau(TS - ST) = \tau((T_{|M}T_0 - T_0T_{|M}) \oplus 0) = \tau(T_{|M}T_0 - T_0T_{|M}) = \mathbf{i}(T_{|M})$, we obtain $ind(T) = \mathbf{i}(T)$.

REMARK 2.6. The trace formula for the index of B-fredholm operators given in Theorem 2.5, extends Fedosov's trace formula for the index of Fredholm operators, see[8, p.10]

3 Properties of the Index

In this section, we will assume that A is a semi-simple complex unital Banach algebra, with unit e, and the ideal J is equal to its socle. Recall that it is well known, that a semi-simple Banach algebra is semi-prime. Then , it follows from [2, BA2.4, p. 103] that an element a in A is invertible modulo J if and only if it is invertible modulo the closure $\overline{J}$ of J. In this case, if p is any minimal idempotent in A, that's a non zero idempotent such that $pAp = \mathbb{C}e$, then the operator $\widehat{a} : Ap \to Ap$, defined by $\widehat{a}(x) = ax$, is a Fredholm operator.
The element $a \in A$ is said to be of finite rank if the operator $\widehat{a}$ is an operator of finite rank. We know from [2, Theorem F.2.4], that the socle of A is $soc(A) =$

$\{x \in A \mid \widehat{x} \text{ is of finite rank}\}$. Moreover, from [1, Section 3], a trace function is defined on the socle by: $\tau(a) = \Sigma_{\lambda \in \sigma(a)} m(\lambda, a)\lambda$, for an element a of the socle of A, where $\sigma(a)$ is the spectrum of a, and $m(\lambda, a)$ is the algebraic multiplicity of λ for a.
For more details about these notions from Fredholm theory in Banach algebras, we refer the reader to [2].
In the following theorem, we will consider stability of B-Fredholmness under small perturbations. Contrarily to the case of usual Fredholmness, we cannot except to preserve B-Fredholmness under perturbation by small norm elements. This can be easily in the case of the algebra $L(X)$. Since 0 is a B-Fredholm operator on the Banach space X, and there exists operators which are not B-Fredholm (see [5, Remark B]), then we cannot have stability of B-Fredholmness under perturbation by small norm elements.

THEOREM 3.1. *Let a be a B-Fredholm element in A modulo J. If $\lambda \in \mathbb{C}, \lambda \neq 0$, and $|\lambda|$ is small enough, then $a - \lambda e$ is a Fredholm element of A modulo J and $\mathbf{i}(a - \lambda e) = \mathbf{i}(a)$.*

Proof. Assume that a is a B-Fredholm element in A modulo J. Then $\pi(a)$ is Drazin invertible in A/J. Let $\Pi : A \to A/\overline{J}$, be the canonical projection. Then $\Pi(a)$ is Drazin invertible in $A/\overline{J}$. As $A/\overline{J}$ is a Banach algebra, then 0 is isolated in the spectrum of $\Pi(a)$ in the Banach algebra $A/\overline{J}$. Thus if $|\lambda|$ is small enough and $\lambda \neq 0$, then $\Pi(a - \lambda e)$ is invertible in $A/\overline{J}$. From our hypothesis on the ideal J, it follows that $\pi(a - \lambda e)$ is invertible in A/J. So $a - \lambda e$ is a Fredholm element in A modulo J.
Moreover,we have from [2, Theorem F.2.6], if p is a minimal idempotent in A, then the operator $\widehat{a - \lambda I} : Ap \to Ap$, defined by $\widehat{a - \lambda I}(x) = (a - \lambda I)x$, is a Fredholm operator on the Banach space Ap. From [10, Theorem 3.17], it follows that $ind(\widehat{a - \lambda I}) = \mathbf{i}(a - \lambda I)$.
On another side, from [3, Remark, iii)], we have $ind(\widehat{a}) = ind(\widehat{a - \lambda I})$, for $|\lambda|$, small enough. Hence $ind(\widehat{a}) = \mathbf{i}(a - \lambda I)$, for $|\lambda|$ small enough.

LEMMA 3.2. *If p is a minimal idempotent in A, then the operator $\widehat{a} : Ap \to Ap$, defined by $\widehat{a}(x) = ax$, is a B-Fredholm operator and $ind(\widehat{a}) = \mathbf{i}(a)$.*

Proof. Since a is a B-Fredholm element in A modulo J, then a is Drazin invertible in A modulo J. From [2, Theorem F.2.4], we know that J is exactly the set of elements x of A such that $\widehat{x}$ is an operator of finite rank. Then $\widehat{a}$ is a Drazin invertible operator modulo the ideal of finite rank on Ap. Thus from Theorem 1.1, $\widehat{a} : Ap \to Ap$ is a B-Fredholm operator. Let $b \in A$ be

a Drazin inverse of a modulo J, then $\widehat{b}$ is a Drazin inverse of $\widehat{a}$ modulo the ideal of finite rank on the Banach space Ap. From Theorem 2.5, we have $\mathbf{i}(\widehat{a}) = \tau(\widehat{a}\widehat{b} - \widehat{b}\widehat{a}) = \tau(\widehat{ab - ba}) = ind(\widehat{a})$. Here τ stands for the trace of the finite rank operators on the Banach space Ap.
Using Lemma 3.2, and since $ab - ba \in J$, it is of finite trace and $\tau(ab - ba) = \tau(\widehat{ab - ba})$. Thus $\mathbf{i}(a) = \mathbf{i}(a - \lambda I)$.

As seen in [4], the product of two B-Fredholm operators in $L(X)$, even it is a B-Fredholm operator, does not have in general its index equal to the sum of the indexes of the operators involved in the product, unless the two operators satisfies a commuting Bezout identity, as proved in [4, Theorem 1.1]. Here we obtain a similar result for the product of B-Fredholm elements.

PROPOSITION 3.3. *Let a_1, a_2 be B-Fredholm elements in A modulo J, and let $\lambda \in \mathbb{C}$.*
i) If $a_1, a_2, u_1, u_2 \in A$ are two by two commuting elements in A such that $u_1a_1 + u_2a_2 = e$, then a_1a_2 is a B-Fredholm element in A modulo J, and $\mathbf{i}(a_1a_2) = \mathbf{i}(a_1) + \mathbf{i}(a_2)$. In particular, if $\lambda \neq 0$, then λa_1 is a B-Fredholm element of A modulo J and $\mathbf{i}(\lambda a_1) = \mathbf{i}(a_1)$.
ii) If j is an element of J, then $a_1 + j$ is a B-Fredholm element in A modulo J and $\mathbf{i}(a_1 + j) = \mathbf{i}(a_1)$.

Proof. i) From [6, Proposition 2.6], it follows that a_1a_2 is a B-Fredholm element in A modulo J. Moreover as $\widehat{u_1}\widehat{a_1} + \widehat{u_2}\widehat{a_2} = I$, and $\widehat{a_1}, \widehat{a_2}, \widehat{u_1}, \widehat{u_2}$, are two by two commuting operators, then from [4, Theorem 1.1], we have $ind(\widehat{a_1}\widehat{a_2}) = ind(\widehat{a_1}) + ind(\widehat{a_2})$, and Lemma 3.2 implies that $\mathbf{i}(\widehat{a_1}\widehat{a_2}) = \mathbf{i}(\widehat{a_1}) + \mathbf{i}(\widehat{a_1})$. Taking $a_2 = \lambda e$, we obtain $\mathbf{i}(\lambda a_1) = \mathbf{i}(a_1)$ because $\mathbf{i}(\widehat{\lambda e}) = ind(\lambda I) = 0$.
ii) If j is an element of J, then $\pi(a_1 + j) = \pi(a_1)$. So $a_1 + j$ is a B-Fredholm element in A modulo J. Moreover if $|\lambda|$ is small enough and $\lambda \neq 0$, then from Theorem 3.1, $a_1 - \lambda e + j$ is a Fredholm element and $\mathbf{i}(a_1 + j) = \mathbf{i}(a_1 - \lambda I + j))$. Using [10, Proposition 3.7, i], we obtain $\mathbf{i}(a_1 - \lambda I + j)) = \mathbf{i}(a_1 - \lambda I) = \mathbf{i}(a_1)$.

References

[1] B. Aupetit, H. du T. Mouton, Trace and determinant in Banach algebras, Studia Math; 121 (1996), 115-136.

[2] B. Barnes, G. J. Murphy, M.R.F. Smyth, T.T. West, Riesz and Fredholm theory in Banach algebras, Pitman Publishing Inc, 1982.

[3] M. Berkani, *On a class of quasi-Fredholm operators*, Integr. Equ. Oper. Theory, 34 (1999), 244-249.

[4] M. Berkani, D. Medkova, A note on the index of B-fredholm operators, Mathematica Bohemica, 129(2004), No.2, 177-180.

[5] M. Berkani, Index of B-Fredholm operators and generalization of a Weyl Theorem, Proc. Amer. Math. Soc., 130 (6) (2002), 1717-1723.

[6] M. Berkani, M. Sarih, An Atkinson type theorem for B-Fredholm operators, *Studia Math.* 148 (2001), 251–257.

[7] M. Berkani, B-Fredholm elements in rings and algebras, In preparation.

[8] A. Böttcher, B. Silbermann, Analysis of Toeplitz operators, Springer, second edition, 2006.

[9] M.D. Cvetkovic, E. Boasso, S.C. Zivkovic-Zlatanovic, Generalized B-Fredholm Banach algebra elements, Mediterranean Journal of Mathematics, pp 1-18, First online: 26 March 2016.

[10] J.J. Grobler, H. Raubenheimer, The index for Fredholm elements in a Banach algebra via a a trace, Studia Mathematica 187(3) (2008), 281-297.

Mohammed Berkani,
Department of Mathematics,
Science faculty of Oujda,
University Mohammed I,
Laboratory LAGA,
Morocco
berkanimo@aim.com,